Assessment of Potential Lake Sturgeon Habitat Availability in Northern Lake Michigan Tributaries: Applications to the Restoration Process

Dan J. Daugherty and Trent M. Sutton
Department of Forestry and Natural Resources
Purdue University
195 Marsteller St.
West Lafayette, IN 47907








- Historic/present status
- Factors
 - Overexploitation
 - Impediments to migration
 - Physical habitat alteration
 - Water-quality degradation




- Increased public awareness, value, appreciation
 - Improved water quality
 - Highly-restricted harvest
- Some stock recovery



- Remaining impediments to recovery
 - Physical habitat alteration
 - Alteration of flow regimes (dams)
 - Land-use practices
 - Erosion/sedimentation
 - Impediments to migration
 - Dams
 - Block access to historic spawning sites
 - Decrease habitat availability
- Relatively permanent, physical changes



- Remediation
 - Rehabilitation of habitats
 - Provide access (fish passage)
 - Removal of barriers
- Current problems
 - Costly processes
 - Limited information exists
 - Where potentially effective?
 - Potential benefit?
 - Which systems?





- Needs:
 - Information on existing habitats
 - Quantification of habitat types
 - Spatial configuration
 - Current accessibility
 - Implementation of restoration strategies
 - Prioritization of systems

Project objectives


- Assess/quantify riverine habitats
 - Egg/larval
 - Juvenile
 - Spawning/staging adult
- Development of digital maps
 - Geo-referenced
 - Spatial relationships
- Develop a habitat-based decision tool
 - Determination of appropriate restoration strategies
 - Prioritization of systems

- Northern Lake Michigan systems:
 - Manistique
 - Fox
 - Ford
 - Menominee
 - Peshtigo
 - Oconto
- Sampled to historic, natural migration barriers


- Suamico
- L. Suamico
- Duck Creek
- Fox




- Stratified, random sampling
 - Run, riffle, pool habitat types
 - Randomly spaced transects
- Point-samples
 - Left, middle, right channel
 - Substrate composition
 - Water depth
 - Flow rate (bottom)
 - Georeferenced (GPS)



- Stratified, random sampling
 - Run, riffle, pool habitat types
 - Randomly spaced transects
- Point-samples
 - Left, middle, right channel
 - Substrate composition
 - Water depth
 - Flow rate (bottom)
 - Georeferenced (GPS)



- Stratified, random sampling
 - Run, riffle, pool habitat types
 - Randomly spaced transects
- Point-samples
 - Left, middle, right channel
 - Substrate composition
 - Water depth
 - Flow rate (bottom)
 - Georeferenced (GPS)





- Additional data considerations
 - Estimation of spring spawning/staging habitat variables
 - Hydrologic discharge/flow relationships
 - Generation of additional seasonal data
 - Seasonal temperature and flow models




Data analyses


- Spatial analyses ArcView/ArcGIS[©]
 - Import coordinates and point-sample data
 - Spatial analyst extension
 - Interpolate variables
 - Create of grids and reclassify data
 - Spatial analyst map calculator
 - Determine habitat availability
 - Known habitat-use characteristics
 - Each life-history stage








Map Calculator



Map Calculator

- Provides estimated feature coverage based on user-specified criteria
- Alter criteria for various life stages and repeat analyses
- Remains geo-referenced





Management benefits

- Realization of enhancement/rehabilitation
- Incorporates all lifestages
- Allows for prioritization of systems
 - Cost-effectiveness
 - Approximation of effort/cost
- Provides framework
 - Behavioral advances
 - Biological advances



Management benefits

- Development of decision tool
 - Applicable to other systems

